Supplementary Table 1. Maternal characteristics (continuous variables).

Variable	Exposed	Unexposed	p-value
Gravida	2 (1–3)	2 (1–3)	>0.05
Parity	2 (1–2)	2 (1–2)	>0.05
Living children	2 (1–2)	2 (1–2)	>0.05
Gestational age (weeks)	38 (37–39)	38 (37–39)	>0.05
Maternal age (years)	30 ± 5	30 ± 5	>0.05

Data are presented as median (IQR) or mean ± SD, as appropriate. SD: Standard deviation.

Supplementary Table 2. Laboratory and low-frequency clinical parameters available in subsets of neonates.

Variable	COVID-19 exposed (n)	Median (IQR)/ mean ± SD	Unexposed (n)	Median (IQR)/ mean ± SD	p-value
C-reactive protein (mg/L)	8	0.13 (0.08–0.36)	2	1.11 (0.69–1.52)	>0.05
Procalcitonin (ng/mL)	8	0.24 (0.15–0.70)	2	0.79 (0.79–0.79)	>0.05
Interleukin-6 (pg/mL)	8	54.85 (21.77–115.26)	2	56.99 (28.56–85.41)	>0.05
Inotrope use (days)	8	0 (0–2)	2	1.5 (0–3)	>0.05
Antibiotic duration (days)	6	7 (7–7)	2	4 (1–6)	>0.05

Data are presented as median (interquartile range) or mean ± SD, as appropriate. These parameters were available only in small subsets of infants and were analyzed descriptively. SD: Standard deviation.