

## A Method to Find out Perceptual Sound Composition of the Vowels in the Contemporary Turkey Turkish

Çağdaş Türkiye Türkçesi Ünlülerinin Algısal Ses İçeriğinin Saptanması için Bir Yöntem

Yusuf Kemal Kemaloğlu<sup>1</sup>, Güven Mengü<sup>2</sup>, Çağıl Gökdoğan<sup>1,3</sup>, Alper Kutalmış Türkcan<sup>1,4</sup>

<sup>1</sup>Gazi University Faculty of Medicine, Department of Otorhinolaryngology and Head-Neck Surgery, Ankara, Turkey

<sup>2</sup>Ankara Hacı Bayram Veli University, Faculty of Letters, Department of Western Languages and Literatures, Ankara, Turkey

<sup>3</sup>Muğla Sıtkı Koçman University Faculty of Medicine, Department of Otorhinolaryngology and Head-Neck Surgery, Muğla, Turkey

<sup>4</sup>University of Health Sciences, Ankara Atatürk Chest Disease and Surgery Training and Research Hospital, Ankara, Turkey

### ABSTRACT

**Objective:** The speech sounds are used in audiology and audio-verbal therapy. Perception of speech sounds is related with their acoustic properties and inner ear physiology. Therefore, a perceptual aspect of the acoustic content should not be overlooked. In this study, we evaluated linear and perceptual changes in the vowels' sound content at and over the comfortable hearing level according to dBA-filter.

**Methods:** Recordings of 8 vowels (<a, e, i, ı, o, ö, u, ü>) of the contemporary Turkey Turkish were filtered by a dBA-filter. Then linear frequency data (Hz) of both original and dBA-filtered files were analyzed for fundamental frequency (F0) and formants (f1 to f5) by Praat; subsequently, the data were transferred to the perceptual range (Critical Bark Bands, CBB).

**Results:** Our data demonstrated that linear values of F0 and f4,5 did not reveal any relationship with vowels, while f1-3 presented phoneme-specific patterns. dBA-filtering did not affect linear data of f3,4 (<u> was the only exception) and f5. Linear f1values were increased by dBA-filter (particularly in <i,u,ü>). f2 of <i, u> presented major deviations. Vowels' CBB-changes were evident in f1 (the only exception was <e>), and it was evident in only f2 of <i, u>.

**Conclusion:** It is apparent that speech sound content at and over the comfortable hearing level stimulates higher frequency bands than found in original voice. Only <e> presented no perceptual change while major changes were particularly seen in <i, u>. Thus, we could pronounce that perceptual aspect by dBA-filter would provide us with a new perspective for understanding the results of speech tests.

**Key Words:** Hearing loss, audiology, speech analysis, vowels, Turkey Turkish

**Received:** 07.16.2018

**Accepted:** 11.04. 2018

### ÖZET

**Amaç:** Konuşma sesleri, odyolojide işitsel değerlendirmede ve işitsel-sözel rehabilitasyonda kullanılmakta olup algılanmaları akustik özellikleri kadar iç kulağın fizyolojisiyle de ilişkilidir. Bu nedenle, akustik içeriğin algısal yönü göz ardı edilmemelidir. Çalışmamızda ünlü fonemlerin doğrusal olan ve olmayan (algısal) ses içeriğinin rahat işitme seviyesinin üzerinde nasıl değiştiğini inceledik.

**Yöntem:** Çağdaş Türkiye Türkçesinin 8 ünlüsü (<a, e, i, ı, o, ö, u, ü>) öncelikle dBA filtresinden geçirildi ve hem orijinal hem de dBA filtresinden geçirilen dosyalar Praat yazılımı kullanılarak doğrusal (Hz) temel frekans (F0) ve formant frekans (f1-f5) değerleri için analiz edildi. Daha sonra doğrusal frekans değerleri algısal değerlere (Kritik Bark Bantları, KBB) çevrildi.

**Bulgular:** F0, f4,5'in doğrusal değerlerinin bütün ünlülerde sabit olduğunu ve sadece f1-3'ün değiştiğini gösterdi. dBA filtresiyle elde edilen sonuçlara bakıldığında; doğrusal f5 değerlerinin değişmediği, f3,4'ün sadece <u>da değişiklik gösterdiği, f1'in doğrusal değerinin ise bütün ünlülerde artarken (özellikle <i, u, ü>de), f2'deki değişim sadece <i, u>da barizdi. Doğrusal olmayan KBB değişimiye, <e> hariç bütün ünlülerde en bariz f1'de görülürken f2 sadece <i, u>da değişmekteydi.

**Sonuç:** Rahat işitme seviyesi ve üzerindeki konuşma seslerinin, orijinal kaydedilen seslere göre daha yüksek frekans bölgelerini uyardığı aşıkardır. Sadece <e> algısal değişiklik göstermezken en büyük değişiklikler <i, u>da görülmektedir. Bu bulgular ışığında, dBA filtresinin algısal frekanslara etkisinin konuşma testlerinin sonuçlarını değerlendirme sürecine yeni bir bakış açısı getirebileceği söylenebilir.

**Anahtar Sözcükler:** İşitme kaybı, odyoloji, konuşma analizi, ünlüler, Türkiye Türkçesi

**Geliş Tarihi:** 16.07.2018

**Kabul Tarihi:** 04.11.201

**ORCID IDs.** Y.K.K. 0000-0001-7236-5576,G.M. 0000-0002-7421-0348,Ç.G.0000-0002-8811-5931,A.T. 0000-0002-9638-455X

**Address for Correspondence / Yazışma Adresi:** Yusuf K. Kemaloğlu, M.D. , Prof. Gazi University Faculty of Medicine, Department of Otorhinolaryngology and Head-Neck Surgery, Besevler, Ankara, Turkey E-mail: yusufk@gazi.edu.tr

©Telif Hakkı 2020 Gazi Üniversitesi Tıp Fakültesi - Makale metnine <http://medicaljournal.gazi.edu.tr/> web adresinden ulaşılabilir.

©Copyright 2020 by Gazi University Medical Faculty - Available on-line at web site <http://medicaljournal.gazi.edu.tr/>

doi:<http://dx.doi.org/10.12996/gmj.2020.91>

## INTRODUCTION

The speech sounds are common test signals for audiological tests and audio-verbal therapy of children with hearing loss (1-6). Although words are mostly used, the isolated vowels and some voiced consonants such as <m, n, s> are also applied as test stimulus during mainly testing young children (Ling sounds) (1,3). For both audiological tests and audio-verbal therapy of children with hearing loss, although the meaning is important, the sound content of the speech sample is essential.

The sound content of the speech samples is composed of complex sound waves. The smallest speech sounds are phonemes or allophones according to causing changes in meaning or not, respectively. There are two kinds of speech sounds in any egressive pulmonary language, vowels, and consonants. Source of vowels' sound energy is vibration of the vocal folds (voicing) while sound source of consonants are either voicing (voiced consonants) or turbulent airflow (unvoiced consonants) which is produced by the specific constriction area (named as articulation region) in the upper airway (particularly in oral cavity) or both (voiced consonants). Sound energy of the vowels and consonants is shaped by filtration and resonance in the anterior portion of the upper airway just after the source and articulation areas, and ultimately a specific sound composition is spread out from the oral and nasal cavities (7).

Under normal conditions, speech sound waves are transferred via the atmosphere and reach the ears of the listeners. Through the external and middle portions of the ear, the sound is transferred to the hairy cells in the Corti organ of the inner ear, in which electrical transformation occurs. Then, the sophisticated sound signal is transferred to the primary hearing cortex, as an electrical stimulus. Although hearing sensation, directly, occurs when the signal is reached to the primary hearing cortex, comprehending of this signal is essential for hearing of speech, which is performed in the secondary and associative hearing and speech areas with contribution of memory regions in the brain (8,9). That means speech comprehension of human beings is based on memorizing language-specific formula of each speech signal. Besides, speech signals are able to carry many other messages to the listeners' brains from the mouths of the speakers, such as information of gender, identity of the speaker, and his or her cultural subgroup and accent (differentiation of regional dialects or whether speakers are native or not) and further feelings, intentions, and metaphors in the mind of the speaker during the speech. In linguistics, all these elements of speech are termed as suprasegmental variables of speech and language (10,11).

It is clear that people are sharing the same language, the native speakers, are able to code all segmental and suprasegmental information via the speech apparatus and decode in the brain.

Details of the speech signal have been subject to many types of research for years. By using different formulas based on fast Fourier transformation (FFT), sound waves in each speech signal have been dissolved (12). As it is known, vowel sounds are produced as nearly periodic complex waves, composed of sound energy accumulated at specific frequency bands, which are known as formants. Consonants are composed of either only aperiodic sound waves or composition of periodic and aperiodic waves (7). For vowels, formants are calculated by using Linear Predictive Coding (LPC) analysis (7,13) or direct visual-manual observation while spectral energy envelopes are produced for the consonants to describe the energy accumulation about frequency ranges (7).

By using the mathematical formulas above, many types of research have been performed to analyze speech sounds. Turkish speech sounds have been first analyzed by Selen (1979) by using a device named as "Sonogramm" (14) and then many researches in which various voice analyzing software were used been done by the linguists, engineers, and physicians (11, 15-19).

In these researches, details of the produced speech sounds have easily been able to be demonstrated. However, as well known in otology and audiology, human hearing is neither one-to-one nor a linear process (20-22). That means sound sample received by the external ear is not transferred to an electrical signal in the Corti organ as it is in details when reached. It has been shown that frequency discrimination in the Corti organ is neither one-for-one nor linear particularly over 500 Hz. The ear is able to hear not each separate frequency but bands and these bands are narrower in low frequencies. That means, the human ear has the ability to distinguish changes in low frequencies better than the high frequencies (21). These frequency bands are known as critical Bark bands (CBB) and the ranges of 24 critical bands were demonstrated between 0 and 18500 Hz by Zwicker & Fastl (1990) (21). Further the following formula (23,24) was also proposed to calculate the Bark values:  $z = (26,81 / (1 + 1960/f)) - 0,53$  ( $z$ : Bark's value;  $f$ : frequency as Hz). The researchers proposed the use of Bark scale to describe the perception of the vowels (21,23-25).

On the other hand, it is known that audibility thresholds of the sounds at different frequencies are different and not linear; the sounds lower than 500 Hz and higher than 6000 Hz can stimulate the Corti organ in higher amplitudes than the sound waves between 500 and 4000 Hz. Furthermore, the external ear canal amplifies sounds particularly at around 2000-3000 Hz; that means, the sounds in this range are perceived as louder than in the atmosphere. dBA filter has been developed to measure the sound energy which harmful to the inner ear, as related with noise issues so that dBA-filter measures the sounds over the audibility thresholds in respect to amplification prior to the inner ear (26,27). Therefore, dBA fits the comfortable hearing level of human hearing at 40 phon as pointed out in the first table of the paper by Barber (2011) (28). British Association of Teachers of the Deaf points out that dBA is used for measures of sound field assessments including speech recognition tests and Assessments with a warble tone generator in Audiology (29).

In this study, as the first and as a preliminary study, we aimed to demonstrate whether the sound content of the vowels in the contemporary Turkey Turkish changed at the comfortable hearing level. These data could help for better understanding of responses of the subjects with hearing loss during audiological tests using the speech data and further audio-verbal therapy.


## MATERIAL and METHOD

This study was performed as a part of TÜBİTAK project, which was designed to develop software to analyze speech and sounds three-dimensionally (3D) in order to produce their 3D printouts. In the project, the voice of a male (age: 43 years) linguist was used for demonstration of analysis and 3D printouts. In this study, we used the recordings to demonstrate a perceptual aspect of the contemporary Turkish vowels. This study was approved by the Ethic Committee for Clinical Studies in Gazi University.

### Testing the dBA filter:

The dBA-filter was run via the software named as "üç boyutlu ses-konuşma analizi programı, 3BKAP" (three-dimensional sound-voice analysis program, figure 1) which was developed as a part of the TÜBİTAK project. This software was based on the Matlab (matrix laboratory), which is computer software to solve mathematical application problems. The following formula used for dBA-filter (<http://www.sengpielaudio.com/BerechnungDerBewertungsfilter.pdf>):

$$A: Ra(f) = \frac{12200^2 \cdot f^4}{(f^2 + 20,6^2) \cdot (f^2 + 12200^2) \cdot \sqrt{f^2 + 107,7^2} \cdot \sqrt{f^2 + 737,9^2}}$$



**Figure 1.** The 3BKAP software used for dBA filtering (arrow 1). Arrow 2 points out the graphic presenting filtered and remained sound content of the file

Before analyzing the data of this study, first, we wanted to demonstrate the output graphics of the dBA-filter in relation to the pure tone signals. For this purpose, first, pure tone sounds (PTS) in the following frequencies were produced in the amplitude of 0,2 Pascal with duration of 0,4 seconds by Praat (30): 125, 250, 500, 1000, 1500, 2000, 3000, 4000, 5000, 6000, 8000, 10000 and 20000 Hz. Then, the PTS were filtered by the 3BKAP, and the PTS and their dBA-filtered outputs were analyzed by the Praat. By using these data, an x-y graphic of the dBA-filter of 3BKAP was formed (x: frequency of the PTAs, y: amplitude, the rest of energy after filtering).

#### Voice recording:

All voice samples of the subject were taken in the silent room of the audiology department by using Shure-SM-58-LCE cardioid microphone with a pop filter concerning Hirano (1989) criteria (31). The Audacity software (2.0.5.) (<https://www.audacityteam.org/>) was used for recordings (sampling rate: 44100 Hz and 16-bit resolution) and saved as wav files. The microphone we used were able to detect the sounds over 50 Hz to 15000 Hz. The subject articulated eight vowels of the contemporary Turkey Turkish, which were symbolized by <a, e, i, i, o, u, ü> in duration as if he was reading aloud the letters in the Turkish alphabet.

#### Quality analysis:

The recorded files were first analyzed by Computerized Speech Laboratory (CSL, Kay Elemetrics, Model, 4300)- "Multidimensional Voice Profile (MDVP)" for mean Schimmer and Jitter values to attain objective quality measures.

#### Filtering:

Then, the recorded (original) files were filtered by dBA-filter and the remaining sound data after filtration was saved as dBA-files of the vowels.

#### Acoustic analysis of the speech sound data:

Both original and dBA-files of the vowels were analyzed for durations, the fundamental frequencies (F0) and formants (f1, f2, f3, f4 and f5) by Praat. The analysis was performed automatically by the script in addition to the Praat version 4.8 (30,32).

By using these data, the following graphics were produced: i) formant graphics, ii) vowel quadrangles by f1 and f2 values; and iii) graphic of intrinsic frequency change (F0 variation vs f1 through the vowels). Then, the CBB values of F0 and all formants were detected according to Traunmüller formula (1988, 1990) (23,24) and the vowel quadrangle was re-formed according to the differences between the CBB values of f2 and f1 (x-axis: f2-f1) and between f1 and F0 (y-axis: f1-F0) as reported by Syrdal & Gopal, (1986) and Traunmüller (1988 and 1990) (23-25).

## RESULTS


Before analyzing the data of the subject, two quality measures were completed. First, the original files produced by the subject of the study were analyzed to demonstrate voice quality of the subject, and it was found that mean Schimmer and Jitter values of the subject were found to be within the normal ranges (Table 1).

**Table 1.** Voice quality during the vowel production by MDVP – CSL

Alphabet symbol	Mean Schimmer	Jitter
<a>	1,09	1,04
<e>	1,56	1,12
<i>	1,24	0,91
<ı>	0,89	0,4
<o>	0,79	4,39
<ö>	0,99	0,34
<u>	0,7	0,45
<ü>	0,64	2,62


Then, the output graphic of the dBA-filter used in this study was prepared and presented in Figure 2. The amplitude of the original PTS was 77 dB-SPL for all frequencies, and after processing by the dBA-filter, the energy lower than 1000

Hz and over 6000 Hz have been decreased while there is an increase from 2000 Hz to 4000 Hz (Figure 2). The amplitude values at 1000 and 6000 Hz by dBA-filter were found to be 77 dB-SPL.



**Figure 2.** dBA filter used in the study (3BKAP software)

After the demonstration of both voice quality of the subject and capacity of the dBA-filter in the software, the study was run by filtering the original files of the eight vowels pronounced by the subject. The graphics seen in Figure 3 reveal the deleted energy and remaining sound contents of the vowels after dBA-filtering. As seen in the figures, the lower frequencies than 1000 Hz lost most of their energy by dBA-filter.



**Figure 3.** Effect of dBA filtering on the original files of the vowels is seen. Blue drawings pointed out the deleted energy by dBA filter; the green drawings are the rest of the speech sample of the vowels

Then, both original and dBA-filtered-files were analyzed by Praat to find out duration, F0 and formant values. As seen in Table 2, dBA-filter did not reveal any significant change in duration. The lowest durations were detected for <i> (0,642 vs 0,641 sec) while the longest ones were for <ö> (0,798 vs 0,797 sec) in both original and dBA files (no statistical difference by Paired sample t-test,  $p > 0,05$ ). Linear (Hz) F0 and formant values revealed by Praat and their CBB values were presented in Tables 3 and 4, respectively.

**Table 2.** Duration of the vowel segments analyzed (seconds)

	original	dBA
<o>	641,60	640,50
<e>	644,56	645,09
<a>	650,50	651,04
<ı>	707,88	708,46
<ü>	714,27	714,25
<i>	773,56	774,20
<u>	795,88	795,86
<ö>	798,03	797,33


**Table 3.** Mean fundamental frequency (F0) and formant frequencies (f1-f5) of the vowels as Hertz (Hz): original files vs dBA files

Alphabet symbols	Files	F0 Mean	f1 Mean	f2 Mean	f3 Mean	f4 Mean	f5 Mean
<a>	original	99,96	707	1104	2773	3283	4050
	dBA	100,41	768	1152	2755	3238	4064
<e>	original	106,44	542	2036	2749	3442	3807
	dBA	110,08	620	2015	2725	3360	3887
<i>	original	104,95	314	2384	2903	3291	3750
	dBA	104,73	462	2368	2870	3306	3778
<ı>	original	103,74	445	1439	2862	3363	4005
	dBA	106,29	738	1800	2972	3430	3966
<o>	original	102,42	593	878	2795	3214	4272
	dBA	102,46	690	910	2800	3222	4227
<ö>	original	97,1	544	1582	2440	3291	4045
	dBA	99,67	649	1598	2443	3282	3991
<u>	original	103,33	349	852	2609	3280	3946
	dBA	102,99	656	1894	2975	3612	3991
<ü>	original	101,95	363	1762	2522	3320	3741
	dBA	103,78	669	1884	2647	3364	3748

**Table 4.** Bark critical band ranges detected for mean fundamental frequency (F0) and formant frequencies (f1-f5) of the vowels according to Trainmüller formula (1988).

Alphabet symbol	Files	Bark for f0 Mean	Bark for F1 mean	Bark for F2 mean	Bark for F3 mean	Bark for F4 Mean	Bark for F5 Mean
<i>	original	1	4	15	16	17	18
	dBA	1	5	15	16	17	18
<ü>	original	1	4	13	15	17	18
	dBA	1	7	13	15	17	18
<e>	original	1	6	14	16	17	18
	dBA	1	6	14	16	17	18
<ö>	original	1	6	12	15	17	18
	dBA	1	7	12	15	17	18
<ı>	original	1	5	11	16	17	18
	dBA	1	7	13	16	17	18
<a>	original	1	7	10	16	17	18
	dBA	1	8	10	16	17	18
<o>	original	1	6	8	16	17	18
	dBA	1	7	8	16	17	18
<u>	original	1	4	8	15	17	18
	dBA	1	7	13	16	17	18

Mean F0 values in the original files presented variations from 97,1 to 106,44 Hz about the phonemes. dBA-filtering did not change F0 in <i, a, o, u>, but caused small increases in the remaining vowels less than 4 Hz (Figure 4).




**Figure 4.** Intrinsic frequency changes (F0 in relation to f1) of the vowels as Hertz (Hz): Original vs dBA files.


There was no correlation between F0 and f1 values in both the original and dBA-files (Paired samples correlation test,  $p > 0,05$ ). The CBB value was found to be 1 for both original and dBA files (Table 3 and Figure 5).

The formant values of the original and dBA files were also seen in Table 3 and Figure 5, and their CBB values were presented in Table 4 and Figure 6. It was apparent that f4 and f5 in both original and dBA files did not present any change in relation to the vowels (figures 5 and 6), while f1, f2, and f3 revealed phoneme-specific patterns.

dBA-filtering did not affect f3 and f4 with the only exception for <u> and f5 in all vowels (Figures 5 and 6). f1 was the major formant affected by dBA-filtering; all f1 values were increased by dBA-filter (up to 1042 Hz in <u>) (Table 3, figure 5) and these increases caused increases in CBB values of the phonemes with the exception of <e>. Major deviations in f1 were seen for <i>, u, ü> while f2 presented major deviations for <i>, u>.




**Figure 5.** Graphics of mean formant frequencies (Hertz, Hz) of the vowels: Original vs dBA files.




**Figure 6.** Graphics of Bark critical band ranges detected for mean fundamental frequency (F0) and formant frequencies (f1-f5) of the vowels according to Trau Müller formula (1988): original files vs dBA files.

In the original files, f2 was closer to f1 in <a>, o, u> vowels and f3 in <i>, ü, e>. Although the distance between f1 and f2 of <a>, o> became closer in dBA-files, major changes in the f1-f2-f3 relationships by dBA-filtering were observed in only <i>, u> (Figure 5). It is clear that in original files, <ö> was the middle vowel of the subject's vowel quadrangle, in which f1 to f2 and f2 to f3 distances were almost equal. By dBA-filtering, <ö> moved to more central position and <i>, u> also became central vowels. In <u>, f2 was increased more than f1 so that the distance between f1 and f2 was increased and f2 became closer to f3.

In the vowel quadrangle, <i>, u, e, a> were the corner vowels of the quadrangle, and <ö, ı> were localized in the center of the quadrangle (Figure 7). By dBA-filtering, all phonemes, but particularly <u, ü, ı>, presented major changes in their locations on the vowel quadrangles. The vowel quadrangle-based on CBB values according to F0-f1 vs. f2-f1 is seen in Figure 8.



**Figure 7.** Vowel quadrangle by linear frequency values (Hertz, Hz): original vs dBA files.



**Figure 8.** Vowel quadrangle by Bark Critical Band values calculated by using Traunmüller formula (1988): original vs dBA files.

## DISCUSSION

In this study, we used the voice of one male subject presenting voice quality data within the normal limits (Table 1). Selen (1979) also reported one male subject's data to describe Turkish phonemes (14). Moreover, Kılıç (2003) and Yılmaz Davutoğlu (2010) reported the data of 5 and four male subjects, respectively, in their studies in which Turkish phonemes were described (16,19). In this study, our purpose was not to describe Turkish vowels acoustically, but to present whether the difference between the sound data produced and perceived at and over the comfortable hearing level (40 phon) could be important regarding audiological tests using speech signals.

Before that, acoustic data found in the original recordings of the subject were reviewed through the literature. It has been reported that the first three formants of the speech are related to phonemic information of the speech (17). In accordance with this assumption, in this study, only  $f_1$ ,  $f_2$  and  $f_3$  revealed differences in relation to the vowels while  $F_0$  and fourth and fifth formants did not present any apparent variation through the phonemes. The subject's Bark Critical Band ranges for  $F_0$ ,  $f_4$  and  $f_5$ , which were 1, 17 and 18, respectively, also supported this notion. Previously, although Kılıç (2003) and Malkoç (2009) reported the graphics using the CBB values of the vowels, they did not present exact CBB values (16,18). However, when we applied Traunmüller (1988) formula (23) to the linear frequency data (Hz) of  $F_0$  and  $f_4$ , which were both reported by Malkoç (2009) (18), it was disclosed that  $F_0$  was 1 and  $f_4$  was 17 (with the exception of <ü> which was 16), as we found. Since the other authors (11,14,16-19) did not present the data of  $f_4$  and  $f_5$  we could only say that CBB value of  $F_0$  was 1 in the studies reporting  $F_0$  (16,18). However, it should be underlined that in this study we used Traunmüller formula (1988, 1990) for CBB conversion of the linear values (Hertz, Hz) (23,24); it is known that the CBB conversion based on the Traunmüller formula (1988, 1990) (23,24) produced some differences in the lower frequencies compared to the original table reported by Zwicker & Fastl (1990) (21). Traunmüller (1990) reported that within the frequency range of the perceptually essential vowel formants (200 – 6700 Hz) the formula agrees to within +/- 0,05 Bark with the Bark scale, originally published in the form of a table (24). Hence, if CBB ranges of  $F_0$  in both our study and the references we pointed out above (16,18) were detected according to the original table of Zwicker & Fastl (1990) (21), it would be 2 for <e, i, o, u, ü> in our study, and for all vowels (<a, e, i, o, ö, u, ü>) in the studies of Kılıç (2003) and Malkoç, (2009) (16,18).

It has been documented that  $f_1$  and  $f_2$  related with amount of the mouth opening and position of the tongue in the anteroposterior direction within the mouth during the articulation of the vowels, respectively because  $f_1$  is a product of the total vocal tract length while  $f_2$  is produced by resonance of the sound within the area in front of the tongue (17). Our  $f_1$  and  $f_2$  values supported the reports of the previous researchers on the Turkish vowels: As demonstrated by the previous researches (14,18,19), <i, ü, e> were the anterior vowels of the contemporary Turkey Turkish in which articulation place was placed anteriorly. Hence, their  $f_2$  was closer to their  $f_3$ , as we found.

On the other hand, <a, o, u> were apparently posterior vowels in which  $f_2$  was closer to  $f_1$  as detected in our study. In this study, the two contradictory vowels (<i, ö>) of the contemporary Turkey Turkish were found to be placed at around the center of the vowel quadrangle, as in accordance with the linear data of Malkoç (2009) (18).

Although <i> as /u/ was demonstrated on the right corner of International Phonetic Association (IPA)'s vowel quadrangle for the contemporary Turkey Turkish (38), Kılıç and Ögüt (2004) revealed that it was a central vowel (39), as we found. The vowel quadrangle-based on  $f_2$ - $f_1$  (x-axis) and  $f_1$ - $F_0$  (y-axis) by using the CBB values supported the data above and it was in accordance with the graphics reported by Kılıç (2003) and Malkoç (2009) (16,18).

IPA documented the following phonetic transcriptions for eight standard vowels of contemporary Turkey Turkish (<a, e, i, o, ö, u, ü>), respectively: /a, e, i, u, o, œ, u, y/ (38). However, Turkish linguistic studies reported more than 16 allophones according to either the acoustic data ( $f_1$  and  $f_2$  values and vowel quadrangle) or perception or sometimes both in addition to various lengthened vowel-uses (10,11,14-16,19,40). When compared linear  $f_1$  and  $f_2$  values (Hertz, Hz) and locations on the vowel quadrangle of our male subject' vowels with the published data about Turkish vowels (11,14-19), it was noticed that our subject produced the following speech sounds for <a, e, i, o, ö, u, ü>, respectively: /a, e, i, ə, o\*, ø, u, y/ (\*the subject pronounced <o> with higher  $f_1$  than reported by Malkoç (2009) and Yılmaz Davutoğlu (2010) (18,19).

Our subject produced vowels within 641 (<o>) to 798 msc (<ö>). The previous papers about Turkish phonemes did not reveal the durations of the vowels that they analyzed. Only studies about the duration of Turkish vowels were done by researchers of the Boğaziçi University (41-43). In their studies performed by using the vowels within the words, they reported the vowel durations for between 49,6 and 184,1 msec. Therefore, since our speaker pronounced them as if he was reading aloud the letters in the Turkish alphabet, our vowels' acoustic data could be considered as examples of the long use of the vowels. Yılmaz Davutoğlu (2010) demonstrated the long allophones of vowels of the contemporary Turkey Turkish as follows (19): /ɑ:/ for <a>, /e:/ for <e>, /i:/ for <i>, /u:/ for <u>, /o:/ for <o>, /ø:/ for <ö>, /y:/ for <y> and /y:/ for <ü>. These demonstrations are in accordance with our data with exception of /ə:/ of <i>. That means, when the based on data of Yılmaz Davutoğlu (which was the only study presenting acoustic details regarding allophones of Turkish vowels) (19), pronunciation of <i> by our subject appears to be artificially related to duration. That is, the contemporary Turkey Turkish includes /ə/ but not its long use /ə:/ in speech according to the data of Yılmaz Davutoğlu (2010) (19).

Behrman (2007) points out an inverse correlation between  $F_0$  and  $f_1$  (intrinsic frequency change) (7) by using the data of Hillenbrand et al. (1995) (33). It is said that  $F_0$  values of <i, u> are expected to be higher than <e, o, a> about increased muscle tension of the tongue on the larynx. Intrinsic frequency change has never been subject to any study performed on the contemporary Turkey Turkish before. In our data, this relationship was partly found;  $F_0$  of <i, u> appeared to be higher than other vowels. When the  $F_0$  and  $f_1$  data reported by Kılıç (2003) and Malkoç (2009) (16,18) were evaluated, intrinsic frequency change was more apparently seen in the data of Kılıç (2003) (16) as reported by Behrman (2007) (7) on the data of Hillenbrand et al. (1995) (33), but partly in the subjects of Malkoç's study (2009) (18), as we found. Further studies are necessary to point out intrinsic frequency changes of the contemporary Turkey Turkish.

Altogether, we could say that our subjects-speaker's vowels are good examples of the contemporary Turkey Turkish. In this study, as the first in the literature, we point out that the sound content of the vowels revealed significant changes when they were filtered by dBA-filter. It is known that dBA roughly corresponds to the inverse of the 40 dB (at 1 kHz) equal-loudness curve for the human ear, which the level of the comfortable hearing (26,28). That means the sound content of the vowels at and over the level of comfortable listening could be different from the produced voice. Since some amount of energy in the lower and upper-frequency ranges are de-emphasized while some in the mid-frequencies between 1000 and 4000 Hz is emphasized, the formant centers and bandwidths should change /shift. It is known that the audiological evaluation which is mostly done at over the comfortable audibility level (2,5,6,29). Therefore, the data derived from the sound content of the produced voice could mislead the clinicians, particularly during fitting the hearing devices and cochlear implants.

Before discussing the differences in the sound content of the dBA-filtered vowels, we should have confirmed that output graphic of our dBA-filter which was prepared as based on the data of the PTS is by dBA filter's graphics reported in the literature. The primary energy which was deleted by this filter was in the lower frequencies than 1000 Hz in PTS data (Figure 2) as well as in the vowels (Figure 3). As a consequence of the energy loss in the lower frequencies in the dBA-files, we noticed that the lowest formant frequency, f1, presented major change and increased. Moreover, the vowel <u> disclosing the lowest formant values among the vowels of the contemporary Turkey Turkish revealed major changes so that its f1, f2, and f3 were all increased. It is noticed that f1 values were increased not only in Hz but also in CBB values.

As seen in the figures 7 and 8, <u> perceptually disclosed a sound composition as in the anterior or middle vowels (<ü, e, ö, >). According to the references about Ling sounds, <u> has been proposed as a signal with lower frequencies; Eastabrooks (2006) reported that f1 and f2 of <u> were 430 and 1170 Hz according to the data of the produced sound content of <u>, respectively (4). The data of this study clearly presented that f1 and f2 values of <u> in our study (349 and 852 Hz, respectively, Table 3) were increased by dBA-filter (656 and 1894 Hz, respectively). Eastabrooks (2006) pointed out the vowels whose f2 was over 1340 Hz as middle and front vowels (4).

Our data reveals that the audible portion of the vowels stimulates higher frequency bands of the Corti than the sound sample found when the produced voiced was analyzed. Therefore, we could suggest that perceptual aspect by using dBA-filter would provide us a new perspective to understand speech tests in the subjects with hearing loss particularly during amplification and audio-verbal therapy. Regarding these aspects, the sound content of the recordings produced by articulation of <e, a, o, i> was appeared to be more convenient than <u, i, ü>. Further studies are necessary to evaluate its clinical importance.

#### Conflict of interest

No conflict of interest was declared by the authors.

#### Acknowledgement

Regarding this study, we would like to express our sincere thanks to The Scientific and Technological Research Council of Turkey (TÜBİTAK) for its great support to our project entitled "Development of a three-dimensional speech analysis program (3BKAP) to form three-dimensional visuals ("phoneme statues") of the phonemes of Turkey Turkish."

#### REFERENCES

- Ling D. Foundations of spoken language for hearing-impaired children. Washington, DC: Alexander Graham Bell Association for the Deaf. 1989.
- Brandy WT. Speech audiometry. In: Katz J, editor. Handbook of Clinical Audiology, Fifth edition. Philadelphia: Lippincott Williams & Wilkins; 2002;96-110.
- Agung K. Bb., Purdy S. C., Kitamura C. The Ling Sound Test Revisited. The Australian and New Zealand Journal of Audiology 2005; 27, 33-41.
- Estabrooks W. Auditory-verbal therapy and practice. In: Estabrooks W, editor. Auditory-verbal therapy and practice. Washington, DC, USA: Alexander Graham Bell Association for the deaf and Hard Hearing, Inc.; 2006; 1-22.
- Güven AG. Bebek ve Çocuklarda İşitimin Davranım Testleriyle Değerlendirilmesi. In: Belgin E, Şahlı S., editörler. Temel Odyoloji. Ankara: Güneş Tıp Kitapevleri; 2014; 231-44.
- Belgin E. Konuşma Odyometresi. In: Belgin E, Şahlı S, editörler. Temel Odyoloji. Ankara: Güneş Tıp Kitapevleri; 2014; 77-82.
- Behrman A. Resonance and articulation. In: Speech and Voice Science. 1<sup>st</sup> ed. San Diego, CA: Plural Pub; 2007;215-351.
- Abbas PJ, Miller CA. Physiology of the Auditory system. In: Cummings CW, Fredrikson JM, Harker LA, Krause CJ, Richardson MA, Schuller DE editörler. Otolaryngology Head & Neck Surgery, Cilt 4. New York: Mosby; 1998, 2831-74.
- Kemaloğlu YK. Sağırın ve sağırlığın nörobiyolojisi. In: Arık E, editör. Ellerle Konuşmak: Türk İşaret Dili Araştırmaları. 1st edition. İstanbul: Koç Ün. Yayınları; 2016. p.86-155. ISBN: 978-605-5250-83-6.
- Özsoy AS. Türkçe'nin Yapısı –I Sesbilim. İstanbul: Boğaziçi Üniversitesi Yayınları; 2004.
- Coşkun V. Türkçenin Ses Bilgisi. İkinci Baskı. İstanbul: IQ Kültür Sanat Yayıncılık; 2010.
- Allen JB, Rabiner LRA. Unified approach to short-time Fourier analysis and synthesis. Proc. IEEE, 1977; 65, 1558-64.
- Kent RD, Read C. The acoustic analysis of speech. San Diego CA: Singular Publishing Group; 1992.
- Selen N. Söyleyiş Sesbilimi, Akustik Sesbilim ve Türkiye Türkçesi, Ankara: Türk Dil Kurumu Yayınları; 1979.
- İclâl E: Türkiye Türkçesinin Görevsel Sesbilimi, Engin Yayıncılık, Ankara,1989.
- Kılıç MA. Türkiye Türkçesindeki ünlülerin sesbilgisel özellikleri. In: Özsoy AS, Akar D, Nakipoğlu Demiralp M, Erguvanlı Taylan EE, Aksu Koç A, editörler. Studies in Turkish linguistics. İstanbul; Boğaziçi Un. Press; 2003, 3-18.
- Türk O, Şaylı Ö, Özsoy AS, Arslan LM. Türkçede ünlülerin formant incelemesi. 18. Dilbilim Kurultayı, 20-21 Mayıs 2004, Ankara.
- Malkoç E. Türkçe ünlü formant frekans değerleri ve bu değerlere dayalı ünlü dörtgeni. Dil Dergisi, 2009; 146: 71-85.
- Yılmaz Davutoğlu A. Standart Türkçedeki ünlülerin akustik analizi ve fonetik altyapı. İstanbul Üniversitesi Sosyal Bilimler Enstitüsü Sahne Sanatları Anasanat Dalı Tiyatro Bölümü. Sanatta Yeterlilik Tezi, İstanbul. 2010.
- Zwicker E, Fastl H. Information processing in the auditory system. In: Psycho-acoustics: Facts and Models. New York: Springer; 1990; 20-55.
- Zwicker E, Fastl H. Critical bands and excitation. In: Psycho-acoustics: Facts and Models. New York: Springer; 1990; 133-55.
- Zwicker E, Fastl H. Just-Noticeable Sound Changes. In: Psycho-acoustics: Facts and Models. New York: Springer; 1990;156-80.
- Traunmüller H. Paralinguistic variation and invariance in the characteristic frequencies of vowels. Phonetica; 1988; 45:1-29.
- Traunmüller H. Analytical expressions for the tonotopic sensory scale. The Journal of the Acoustical Society of America; 1990; 88: 97.
- Syrđal AK, Gopal HS. A perceptual model of vowel recognition based on the auditory representation of American English vowels. Journal of Acoustical Society of America; 1986;79:1086-100.
- Levey S, Fligor BJ, Ginocchi C, Kagimbi L. The effects of noise-induced hearing loss on children and young adults. Contemporary Issues in Communication Science and Disorders; 2012; 39:76-83.
- Kemaloğlu YK, Tutar H. Gürültüye bağlı işitme kayıpları ve akustik travma. Türkiye Klinikleri J E.N.T.-Special Topics; 2013, 6(1), 44-54
- Barber JR, Warner KA, Theobald DM. Anthropogenic noise exposure in protected natural areas: estimating the scale of ecological consequences. Landscape Ecology; 2011; 26:1281-95.
- Westhorp S. Decibel Scales in Audiology In: Underwood A., editor, Audiology refreshers. BATOD, 2009. <https://www.batod.org.uk/wp-content/uploads/2018/02/Audiology-Refreshers.pdf>
- Boersma P, Weenink D. Praat: doing phonetics by computer. [Bilgisayar programı]. İndirilme adresi: <http://www.praat.org/>, 2013.
- Hirano M. Objective evaluation of the human voice: clinical aspects. Folia Phoniatr (Basel); 1989;41:89-144.
- Kılıç MA. Özelleştirilmiş Praat Programı ile Ses, Rezonans ve Konuşma Değerlendirmesi Kursu, 10. Ulusal Larengoloji Kongresi, 4-5 Mayıs 2018, Ankara.
- Hillenbrand J, Getty LA, Clark MJ, Wheeler K. Acoustic characteristics of American English vowels. Journal of Acoustical Society of America; 1995, 97:3099-3111.
- Göksel A, Kerslake C. Phonologic units. In: Turkish: A Comprehensive grammar. London: Routledge; 2005; 3-13.
- Şaylı Ö, Levent M, Arslan A, Özsoy S. Duration Properties of the Turkish Phonemes, 11th International Conference on Turkish Linguistics, Gazimağusa, KKTC, Ağustos 7-9 2002.
- Şaylı Ö, Arslan LM. Türkçe'deki seslerin süre özellikleri. Dilbilim Araştırmaları, Boğaziçi University Press; 2003; 15-26,
- Arisoy E, Arslan LM, Demiralp MN, Ekenel HK, Kelepir M, Meral HM, et al. Duration of Turkish Vowels Revisited. 2004.
- Zimmer K, Orgun O. Turkish. In: The handbook of International Phonetic Association. Chambridge: Chambridge University Press; 1999, 154-8.
- Kılıç MA, Öğüt F. A high unrounded vowel in Turkish: is it a central or back vowel? Speech Communication; 2004, 43: 143-154.
- Göksel A, Kerslake C. Phonologic units. In: Turkish: A Comprehensive grammar. London: Routledge; 2005, 3-13.
- Şaylı Ö, Levent M, Arslan A, Özsoy S. Duration Properties of the Turkish Phonemes, 11th International Conference on Turkish Linguistics, Gazimağusa, KKTC, Ağustos 7-9 2002.
- Şaylı Ö, Arslan LM. Türkçe'deki seslerin süre özellikleri. Dilbilim Araştırmaları, Boğaziçi University Press; 2003; 15-26,
- Arisoy E, Arslan LM, Demiralp MN, Ekenel HK, Kelepir M, Meral HM, et al. Duration of Turkish Vowels Revisited. 2004.